



RED FLAGS

Síndrome de Sjögren



FACTORES PREDICTORES DE LINFOMA

SÍNDROMES LINFOPROLIFERATIVOS



- Hombre
- Parotidomegalia
- Crioglobulinas +
- Hipocomplementemia
- Edad de debut <35 años
- Adenopatías
- Anemia, trombopenia, neutropenia
- Pico monoclonal
- Factor Reumatoide +

- Linfoma MALT
- Linfoma B difuso de células grandes
- Linfoma nodal de la zona marginal



No olvidar mantener vigilancia activa para otros tumores:
-Páncreas
-Tiroides
-Pulmón

Bibliografía:

1. Brito-Zerón P, Kostov B, Fraile G, Caravia-Durán D, Maure B, Rascón FJ, Zamora M, Casanovas A, Lopez-Dupla M, Ripoll M, Pinilla B, Fonseca E, Akasbi M, de la Red G, Duarte-Millán MA, Fanlo P, Guisado-Vasco P, Pérez-Alvarez R, Chamorro AJ, Morcillo C, Jiménez-Heredia I, Sánchez-Berná I, López-Guillermo A, Ramos-Casals M; SS Study Group GEAS-SEMI. Characterization and risk estimate of cancer in patients with primary Sjögren syndrome. *J Hematol Oncol.* 2017 Apr 17;10(1):90. doi: 10.1186/s13045-017-0464-5. PMID: 28416003; PMCID: PMC5392920.
2. Paul Creamer, MD. Samuel Moschella, MD. Clinical manifestations of Sjögren's disease: Extraglandular disease. In: UpToDate (Accessed on August 18, 2023)